

## **DUFA-START**

### **Composition:**

Contains per litre solution:

|                                   |          |
|-----------------------------------|----------|
| Magnesium sulphate .....          | 0.91 g   |
| Citric acid .....                 | 6.56 g   |
| Sodium chloride .....             | 19.63 g  |
| Potassium chloride.....           | 164.85 g |
| Riboflavin sodium phosphate ..... | 0.07 g   |

### **Indications:**

Prevention of dehydration and loss of vitality in poultry in periods of stress (disease, vaccination, high ambient temperatures, rapid weight gains) by stimulation of water consumption and feed intake.

### **Contraindications:**

None.

### **Dosage and administration:**

For oral administration in drinking water.

1 litre per 500 litres of drinking water.

Dose per 1000 animals per day per age category:

Broilers:

|                     |        |
|---------------------|--------|
| 1 week .....        | 30 ml  |
| 2 weeks.....        | 60 ml  |
| 3 weeks.....        | 90 ml  |
| 4 weeks and up..... | 120 ml |

Layers and breeding stock:

|                      |            |
|----------------------|------------|
| 1-5 weeks .....      | 30 ml      |
| 6-10 weeks .....     | 50 ml      |
| 11-15 weeks .....    | 80 ml      |
| 16 weeks and up..... | 120-200 ml |

Shake well before use.

Once diluted in water Dufa-Start should be used immediately.

### **Withdrawal times:**

None.

### **Storage conditions:**

Store dry and below 25°C. Do not freeze.

### **Packing:**

1 litre bottles and 5 litre cans.

