

DUFACOX 2.5% ORAL

Composition:

Contains per ml solution:

Toltrazuril25 mg

Description:

Dufacox 2.5% Oral is a anticoccidial which is active against all intracellular development stages of coccidia. Toltrazuril is effective in the treatment of coccidiosis in poultry, turkeys, calves, lambs, kids and piglets caused by different species of *Eimeria* spp.

Indications:

Coccidiosis of all stages like schizogony and gametogony stages of different *Eimeria* spp. in chickens, turkeys, calves, lambs, kids and piglets.

Contraindications:

Do not administer to animal with an impaired liver and/or renal function. Do not use in breeders in production or layers producing eggs for human consumption.

Side effects:

At high dosages in laying hens egg-drop and in broilers growth inhibition and polyneuritis can occur.

Dosage and administration:

For oral administration via drinking water.

Poultry and turkeys: 1 litre per 1000 litres of drinking water continuous medication for 2 days (7.5 mg/kg body weight)

Calves, lambs, kids and piglets: 0.8 ml per kg body weight for 2 days (20 mg/kg body weight).

During treatment the medicated water should be the only source of drinking water!

Medicated water should be used immediately and refreshed every 24 hours for the duration of the treatment.

Do not use in layers producing eggs for human consumption.

Withdrawal times:

Meat: 19 days (poultry and turkeys)

42 days (sheep and goats)

63 days (calves)

77 days (pigs)

Storage conditions:

Store dark and below 25°C. Do not freeze.

Packing:

1 litre cans.

