

DUFAMEC 1.02% ORAL

Composition:

Contains per ml oral solution:

Ivermectin 10.2 mg

Description:

Oral solution containing ivermectin, a broad spectrum anthelmintic for the control of a wide variety of endo- and ectoparasites.

Indications:

Endo- and ectoparasitic infections in horses, sheep, goats and poultry, caused by parasites susceptible to ivermectin.

Horses: *Strongylus vulgaris* (adult and arterial larval stages), *S. edentatus* (adult and tissue stages), *S. equinus* (adults), *Triodontophorus* spp. (adults), *Cyathostomum*, *Cylicocylus* spp., *Cylicostephanus* spp., *Cylicodontophorus* spp., *Gyalocephalus* spp. (adults and non-inhibited mucosal larvae), *Parascaris equorum* (adults and larvae), *Oxyuris equi* (larvae), *T. axei* (adults), *Strongyloides westeri* (adults), *Habronema* spp. (adults), *Onchocerca* spp. (microfilariae), *Dictyocaulus arnfieldi* (adults and larvae), *Gastrophilus* spp. (larvae)

Sheep: *Haemonchus contortus*, *Ostertagia circumcincta*, *Trichostrongylus* spp., *Cooperia* spp., *Nematodirus* spp. including *N. battus*, *Strongyloides papillosus*, *Oesophagostomum* spp. and adult *Charbertia ovina*. Inhibited larval stages and benzimidazole resistant strains of *H. contortus* and *O. circumcincta* are also controlled; *Dictyocaulus filaria*, *Oestrus ovis*.

Goats: *H. contortus*, *O. circumcincta*, *T. colubriformis*, *N. spathiger*, *Strongyloides papillosus*, *O. columbianum* and adult *Charbertia ovina*, *Dictyocaulus filaria*.

Poultry: *Ascaridia* spp., *Capillaria* spp., *Cnemidocoptes mutans* and subcutaneous mite, lice infestations.

Contraindications:

Hypersensitivity to ivermectin.

Dosage and administration:

For oral administration.

The recommended dose is 0.2 mg ivermectin per kg body weight.

Horses 1 ml per 50 kg body weight

Sheep and goats 0.2 ml per 10 kg body weight

In sheep and goats, administer with a dosing gun. The dosing gun should be calibrated accurately to ensure precise dosing of young sheep and goats.

Poultry 40 ml per 250 – 300 litres of water (1000 birds)

Shake well before use.

Side effects:

Hypersensitivity symptoms caused by immunological reactions to dying or dead microfilariae and migrating larvae can occur.

Precautions and warnings:

Dose accurately to prevent development of resistance caused by underdosing.

Withdrawal times:

Meat: 35 days (horses)

14 days (sheep, goats, poultry)

Eggs: 10 days

Do not use in animals producing milk for human consumption.

Storage conditions:

Store dark and below 25°C. Do not freeze.

Packing:

100 ml and 1 litre bottles.