

DUFAMOX-C 200/2 mln W.S.P.

Composition:

Contains per g water soluble powder:

Amoxicillin trihydrate 200 mg
Colistin sulphate.....2,000,000 IU

Description:

Amoxicillin is a semisynthetic beta-lactam antibiotic effective against a broad range of gram-negative and gram-positive bacteria. Amoxicillin is susceptible to degradation by β -lactamase-producing bacteria.

Colistin is a polymyxin antibiotic with bactericidal activity against certain gram-negative bacteria, including *Salmonella* spp. and *E. coli*. Absorption of colistin sulphate after oral administration is minimal and it is therefore only active in the gastrointestinal tract. In addition to the direct antibacterial activity, colistin also exhibits potent anti-endotoxin activity.

Combined, amoxicillin and colistin act synergistically.

Indications:

Gastrointestinal, respiratory and urinary tract infections caused by microorganisms susceptible to amoxicillin and colistin.

Contraindications:

Hypersensitivity to penicillins or colistin, severe renal impairment.

Dosage and administration:

For oral administration in drinking water

Recommended dose 500 g per 1000 litres drinking water for 3-5 days

Medicated water should be used immediately and refreshed every 24 hours for the duration of the treatment.

Side effects: gastrointestinal disturbances.

Precautions and warnings:

It is recommended that the susceptibility of the causative microorganisms is established before treatment is commenced. Severely sick animals have altered drinking patterns and may need to be medicated parenterally. Dosage may need to be adjusted to reflect herd or flock water consumption. Do not handle the product if you are known to be hypersensitive to penicillins. If hypersensitivity reactions occur, seek immediate medical advice.

Withdrawal times:

Meat: 8 days

Eggs: 15 days

Storage conditions:

Store dry and below 25°C. Do not freeze.

Packing:

500 g and 1 kg jars.

