

DUFANAZEN

Composition:

2.36 g contains:

Diminazene diacetate 1.05 g

Antipyrine 1.31 g

Description:

Diminazene diacetate is an antibacterial and antiprotozoal active against babesia, piroplasmosis and trypanosomiasis. Diminazene diacetate is indicated for prophylaxis and treatment of babesia, piroplasmosis and trypanosomiasis in camel, cattle, cats, dogs, goats, horse, sheep and swine. Antipyrine, also known as phenazone, is an analgesic and antipyretic.

Indications:

Treatment and prevention of protozoan diseases in cattle, sheep, goats, horses and dogs.

- Trypanosomosis caused by *Trypanosoma congolense*, *T. vivax*, *T. brucei*, *T. evansi* and *T. equiperdum*.
- Piroplasmosis caused by *Babesia bovis*, *B. bigemina*, *B. ovis*, *B. motasi*, *B. divergens*, *B. caballi*, *B. equi*, *B. gibsoni* and *B. canis*
- Theileriosis caused by *Theileria annulate*.

Contraindications:

Hypersensitivity to any of the active ingredients. Do not administer in animals with impaired renal and/or hepatic functions. Theileriosis can interfere with the immune system, therefore a vaccination program should be postponed.

Dosage and administration:

For deep intramuscular administration.

Reconstitute solution by dissolving 2.36 g of Dufanazen in 15 ml sterile water to obtain a ready-to-use 7% diminazene diacetate solution.

Treatment of piroplasmosis and trypanosomosis caused by *T. Congolense* and *T. vivax*:

1 ml per 20 kg body weight (2.36 g of Dufanazen is sufficient for 300 kg)

Treatment of trypanosomosis caused by *T. brucei* and theileriosis caused by *Theileria annulate*:

1 ml per 10 kg body weight (2.36 g of Dufanazen is sufficient for 150 kg)

Do not exceed a total dose of 4 g of active ingredient in any animal.

If the animal does not recover after treatment consult a veterinarian.

Divide large volumes injections over 2 separate sites.

Inject dogs strictly according to body mass. Overdosage or repeated dosing may cause central nervous symptoms in dogs.

Withdrawal times:

Meat and offal: 21 days

Milk: 3 days

Storage conditions:

Store dark, dry and below 25°C. Do not freeze.

The reconstituted solution can be stored during 15 days in a fridge, if protected from light and in a closed glass vial.

Packing:

2.36 g sachet, 150 sachets per carton;

23.6 g sachet, 10 sachets per carton.