

DUFAPIRIN C+K3 W.S.P.

Composition:

Contains per g watersoluble powder:

Sodium salicylate	300 mg
Ascorbic acid (C).....	250 mg
Menadione sodium bisulfite (K3)	25 mg

Description:

Sodium salicylate is an anti-inflammatory, antipyretic and analgesic agent belonging to the non-steroidal anti-inflammatory drugs group (NSAIDs).

Vitamin C (ascorbic acid) is essential for the formation and maintenance of the normal function of skeletal tissues and exerts an important influence on the reproductive system.

Vitamin K3 (as menadione sodium bisulfite) plays an important role in the coagulation process of blood and also in the cell respiration process.

Indications:

For the treatment of pain, inflammation and pyrexia as anti-pyretic, analgesic, anti-inflammatory, antiphlogistic and platelet anti-aggregating in viral infections, acute bacterial processes or infective syndromes. For osteoarticular and muscular disorders, rheumatoid episodes, fever syndromes, heating stress in calves, swine and broilers.

Contraindications:

Do not use in suckling animals or dehydrated animals. Do not use in animals with digestive tract disorders, especially in case of gastro-intestinal ulcerations and/or haemorrhage and neither in case of pre-existing cardiac, renal or hepatic disease. Do not use in case of hypersensitivity to the active substances. It can provoke stomach injuries especially in sensitive animals.

Dosage and administration:

For oral administration in drinking water.

Broilers: 0.1 g per kg body weight, or 100 g per 200 litres of drinking water for 1000 birds, per day, for maximum 5 days.

Calves and swine: 0.1 g per kg body weight, per day, for maximum 5 days.

Medicated water should be used immediately and refreshed every 24 hours for the duration of the treatment. The body weight should be determined with accuracy.

Precautions and warnings:

Do not mix with feed. Avoid combination with other non-steroidal anti-inflammatory drugs and with amino-glycoside antibiotics. Do not mix with other veterinary medicinal products. Do not exceed the recommended dosage levels. For pregnancy, lactation, laying: use only according to the benefit/risk assessment by the responsible veterinarian.

Withdrawal times:

None.

Do not use in animals producing milk or eggs for human consumption.

Storage conditions:

Store dry and below 25°C. Do not freeze.

Packing:

1 kg jars.