

LEVORAL 75% W.S.P.

Composition:

Contains per g water soluble powder:

Levamisole (as hydrochloride)..... 750 mg

Description:

Levamisole is an imidazothiazole anthelmintic and elicits spastic muscle paralysis in the parasite. Its antinematodal activity is effective against lung and gastro-intestinal nematodes.

Indications:

Endoparasitic infections in cattle, sheep, swine and poultry, caused by parasites susceptible to levamisole. Cattle and Sheep: *Haemonchus*, *Trichostrongylus*, *Ostertagia*, *Cooperia*, *Nematodirus*, *Bunostomum*, *Oesophagostomum*, *Chabertia*, and *Dictyocaulus* (lungworm) spp.. Swine: *Metastrongylus elongatus* (lungworm) and the gastrointestinal nematodes *Ascaris suum* and larva migrans, *Hyoststrongylus rubidis*, *Strongyloides* spp., *Oesophagostomum dentatum* and *Trichuris suis*. Poultry: *Ascaridata* and *Capillaria* spp.

Contraindications:

Hypersensitivity to levamisole.

Dosage and administration:

For oral administration in drinking water.

Cattle, sheep and swine..... 1 g per 100 kg body weight

Poultry..... 15 g per 100 litres of water

Medicated water should be used immediately.

Side effects:

At recommended doses levamisole rarely causes side effects. At higher doses transient symptoms may appear, these include head shaking, salivation and slight muscle tremors. These are more likely to occur in cattle than sheep. Shock has been observed in animals with serious and heavy worm infections after repeated treatment.

Interactions:

Do not combine with other drugs with a 'nicotine-like' action (e.g., pyrantel and morantel).

Precautions and warnings:

The use of disposable gloves and facemasks during preparation is recommended.
Dose accurately to prevent development of resistance caused by underdosage.

Withdrawal times:

Meat: 7 days; eggs: 3 days.

Storage conditions:

Store below 25°C. Do not freeze.

Packing:

100 sachets and 1 kg jar.

