

NICLO-LEVA TABLETS

Composition:

Contains per tablet:

Niclosamide 450 mg

Levamisole HCl 50 mg

Metoclopramide 2.5 mg

Description:

Effective broad spectrum wormer for the control of roundworms, hookworms and tapeworms (except Hydatids) in dogs and cats.

Indications:

Roundworms (*Toxocara* spp and *Toxocaris leonina*), hookworms (*Unicinaria stenocephala* and *Ancylostoma* spp), and tape worms (*Dipylidium* spp and *Taenia* spp but not *Echinococcus granulosus* or Hydatids).

Dosage and administration:

Recommended dose 1 tablet per 5 kg body weight

Give tablet before or with a small meal.

It is recommended that tablets be crushed and added to food.

Treat adult animals at least twice, preferably four times a year.

Breeding animals should be treated twice during pregnancy at a 14 day interval and every three to four weeks during lactation.

Treat puppies and kittens every fortnight until 12 weeks of age.

For kittens and old cats divide the dose giving 1/2 the dose after a morning meal and the same dose 8 – 12 hours later.

Side-effects:

Do not overdose. Overdosage may cause excess salivation, vomiting and disorientation.

Gastrointestinal upset may occur in rare cases.

Storage conditions:

Store dry and below 25°C. Do not freeze.

Packing:

Jar containing 500 tablets.