

## **THIAMPHENICOL 10% ORAL**

### **Composition:**

Contains per ml oral solution:

Thiamphenicol..... 100 mg

### **Description:**

Thiamphenicol is a broad-spectrum antibiotic with bacteriostatic activity against gram-positive and gram-negative bacteria, in particular *Spirochetes*, *Rickettsiae*, *Pasteurella*, *Treponema* and *Bordetella* spp.

### **Indications:**

Gastrointestinal and respiratory infections in cattle, swine and poultry, caused by microorganisms susceptible to thiamphenicol.

### **Contraindications:**

Hypersensitivity to thiamphenicol. Animals with any evidence of pre-existing bone-marrow depression or blood dyscrasias.

### **Dosage and administration:**

For oral administration in drinking water.

Therapeutic dose ..... 1 ml per litre of water

Prophylactic dose.....0.5 ml per litre of water

Shake well before use. Medicated water should be used immediately and refreshed every 24 hours for the duration of the treatment.

### **Side effects:**

Dose-dependent reversible depression of the bone-marrow, haematological dyscrasias, gastrointestinal disturbances and cutaneous eruptions.

### **Interactions:**

Do not combine with bactericidal antibiotics or sulfamethoxypyridazine.

### **Precautions and warnings:**

It is recommended that the susceptibility of the causative microorganisms is established before treatment is commenced.

Severely sick animals have altered drinking patterns and may need to be medicated parenterally.

Dosage may need to be adjusted to reflect herd or flock water consumption.

Use in pregnant or lactating animals only after risk/benefit assessment by veterinarian.

Reduce dosage in animals with impaired kidney function.

### **Withdrawal times:**

Meat: 18 days

**Storage conditions:**

Store below 25°C. Do not freeze.

**Packing:**

100 ml and 1 litre bottles.