

TRIM/SUL D 40/200 INJ

Composition:

Contains per ml solution:

Trimethoprim..... 40 mg
Sulfadoxine 200 mg

Description:

Trimethoprim is an antibacterial agent with a bacteriostatic activity against many gram-negative and gram-positive bacteria. Sulfadoxine is a sulfonamide antibiotic with bacteriostatic activity. The combination ('potentiated sulphas') is bactericidal against a wide variety of gram-positive and gram-negative bacteria as trimethoprim and sulfadoxine act synergistically.

Indications:

Gastrointestinal, respiratory and urogenital infections in cattle, swine, sheep, goats and horses caused by microorganisms susceptible to the combined effects of trimethoprim and sulfadoxine, such as acute or chronic enteritis caused by *Salmonella choleraesuis* and some other *Salmonella* spp.; post-weaning diarrhoea caused by K88-positive, K99-positive or 987P *Escherichia coli*-strains; secondary bacterial infections with *Pasteurella multocida*, *Actinobacillus pleuropneumoniae*, *Streptococcus* spp and *Haemophilus parasuis*, during the course of viral disease.

Contraindications:

Hypersensitivity to trimethoprim or sulphonamides, blood dyscrasias or severe hepatic or renal impairment.

Dosage and administration:

For intramuscular (cattle, swine, sheep, goats) or slow intravenous (cattle, horse) administration. Recommended dose 1 ml per 10 kg body weight per day for up to 5 days

Side effects:

Transient injection site reactions.

Interactions:

Do not administer concurrently with members of the Vitamin B complex, or local anaesthetics such as procaine and lidocaine.

Withdrawal times:

Meat: 14 days
Milk: 4 days

Storage conditions:

Store dark and below 25°C. Do not freeze.

Packing: 100 ml vials.

